



Aqueous two-phase systems for extractive enzymatic hydrolysis of biomass

Bussamra, Bianca Consorti; Azzoni, Sindelia Freitas; Mussatto, Solange I.; da Costa, Aline Carvalho; van der Wielen, Luuk A. M.; Ottens, Marcel

Publication date:
2017

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Bussamra, B. C., Azzoni, S. F., Mussatto, S. I., da Costa, A. C., van der Wielen, L. A. M., & Ottens, M. (2017). *Aqueous two-phase systems for extractive enzymatic hydrolysis of biomass*. Abstract from Biopartitioning and Purification Conference 2017, Copenhagen, Denmark.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

AQUEOUS TWO-PHASE SYSTEMS FOR EXTRACTIVE ENZYMATIC HYDROLYSIS OF BIOMASS

Bianca Consorti Bussamra^{1,2}, Sindelia Freitas Azzoni², Solange I. Mussatto³, Aline Carvalho da Costa², Luuk van der Wielen¹, Marcel Ottens^{*1}

¹*Delft University of Technology. Van der Maasweg 9, 2629HZ. Delft, The Netherlands.*

²*University of Campinas. Av. Albert Einstein, 500. Post Code: 6066. Campinas, Brazil.*

³*Technical University of Denmark. Kemitorvet, Building 220, 2800. Kongens Lyngby, Denmark.*

Sugars derived from lignocellulosic materials are the main carbon sources in bio-based processes aiming to produce renewable fuels and chemicals. One of the major drawbacks during enzymatic hydrolysis of lignocellulosic materials to obtain sugars is the inhibition of enzymes by reaction products (cellobiose and glucose). This effect is even more pronounced in hydrolysis containing high solid content (15-20% or higher water-insoluble solids – WIS), which is desired in order to obtain hydrolysates containing high total reducing sugar concentration and reduced water usage¹. The aim of this project is to develop a new process for sugarcane bagasse hydrolysis using aqueous two-phase system. This system will be applied as *in situ* extraction aiming to remove the reaction products as they are released. As a consequence of product removal, enzymes tend to maintain their maximum activity². The phase-components of the systems will be chosen taking into account their costs, viscosity, capacity of regeneration, melting point, solubility and partition of sugars and proteins. The pre-selected components will be studied and tested in high-throughput experiments³, in order to determine their partition coefficients of sugars and enzymes, phase diagrams and volumetric ratios. The results of this project will make possible to design a process that enables high sugar concentration during the hydrolysis reaction, overcoming one of the biggest drawbacks regarding the production of second-generation ethanol: the enzymatic inhibition. The achievement of the project's goal can lead to, but not limited to, three consequences: enhancement of sugarcane mills productivity; reduction of fossil fuels usage, which can accelerate the energetic independence in many countries; and contribution to a more sustainable economy. This paper will present optimal aqueous two-phase systems for the separation of sugars and enzymes, which allow the development of an improved second-generation ethanol process.


References

1. Modenbach, A. A. & Nokes, S. E. (2013) Enzymatic hydrolysis of biomass at high-solids loadings - A review. *Biomass and Bioenergy* 56: 526–544.
2. Yang, J., Zhang, X., Yong, Q. & Yu, S. (2011). Three-stage enzymatic hydrolysis of steam-exploded corn stover at high substrate concentration. *Bioresour. Technol.* 102: 4905–4908
3. Bensch, M., Selbach, B. & Hubbuch, J. (2007) High throughput screening techniques in downstream processing: Preparation, characterization and optimization of aqueous two-phase systems. *Chem. Eng. Sci.* 62: 2011–2021.



Biopartitioning and Purification Conference 2017

SEPTEMBER 24-27 | COPENHAGEN, DENMARK | LUND UNIVERSITY



Posters:**(P1) DESIGN AND OPERATION OF A CONTINUOUS SEPARATOR FOR AQUEOUS TWO-PHASE SYSTEMS**

B.A. Andrews[#], E.C. Huenupí and J.A. Asenjo

Centre for Biotechnology and Bioengineering .Department of Chemical Engineering and Biotechnology, University of Chile, Santiago, Chile.

(P2) AQUEOUS TWO-PHASE SYSTEMS FOR EXTRACTIVE ENZYMATIC HYDROLYSIS OF BIOMASS

Bianca Consorti Bussamra^{1,2}, Sindelia Freitas Azzoni², Solange I. Mussatto³, Aline Carvalho da Costa², Luuk van der Wielen¹, Marcel Ottens^{*1}

¹*Delft University of Technology. Van der Maasweg 9, 2629HZ. Delft, The Netherlands.*

²*University of Campinas. Av. Albert Einstein, 500. Post Code: 6066. Campinas, Brazil.*

³*Technical University of Denmark. Kemitorvet, Building 220, 2800. Kongens Lyngby, Denmark.*

(P3) ENHANCED CATALYTIC ACTIVITY AND STABILITY OF CYTOCHROME C IN BIO-IONIC LIQUIDS

Meena Bisht,^a Dibyendu Mondal,^b Matheus M. Pereira,^b Mara G. Freire,^b P. Venkatesu,^{a*} and J. A. P. Coutinho^{b*}

^a*Department of Chemistry, University of Delhi, Delhi 110 007, India*

^b*Departamento de Química, CICECO, Universidade de Aveiro, Aveiro, Portugal*

(P4) PARTITION BEHAVIOR OF LACCASE IN IONIC-LIQUID-BASED AQUEOUS BIPHASIC SYSTEMS

Ana P.M. Tavares^{1*}, Oscar Rodriguez², João A.P. Coutinho¹, Ana Soto², Mara G. Freire¹

¹*CICECO-Aveiro Institute of Materials, Chemistry Department, University of Aveiro, 3810-193 Aveiro, Portugal, e-mail: aptavares@ua.pt*

²*Department of Chemical Engineering, Universidade de Santiago de Compostela, Santiago de Compostela, Spain*

(P5) THE HYDROGEN BOND BASICITY OF IONIC LIQUIDS AS A TOOL FOR PREDICTING THE FORMATION OF AQUEOUS BIPHASIC SYSTEMS

Helena Passos¹, Teresa B. V. Dinis¹, Ana Filipa M. Cláudio¹, Mara G. Freire¹ and João A. P. Coutinho^{1,*}

¹*CICECO - Aveiro Institute of Materials, Department of Chemistry, University of Aveiro, Aveiro, Portugal*

(P6) FACTORS DRIVING SOLUTE PARTITION IN IONIC-LIQUID-BASED AQUEOUS BIPHASIC SYSTEMS

Helena Passos¹, Teresa B. V. Dinis¹, Emanuel V. Capela¹, Maria V. Quental¹, Joana M. Gomes¹, Judite Resende¹, Pedro P. Madeira¹, Mara G. Freire¹ and João A. P. Coutinho^{1,*}

¹ *CICECO - Aveiro Institute of Materials, Department of Chemistry, University of Aveiro, Aveiro, Portugal*

ADSORPTION OF NON-VOLATILES FROM FOOD PRODUCTS

Mónica Moreno-González^a, Nienke Hylkema^b, Guilherme Ferreira^b, Hilde Wijngaard^c, Marcel Ottens^{a,*}

^a*Department of Biotechnology, Delft University of Technology, Van der Maasweg 9, 2629HZ Delft, The Netherlands*

^b*DSM Food Specialties, Alexander Flemminglaan1, 2613 AX Delft, The Netherlands*

^c*Unilever R&D Vlaardingen, Olivier van Noortlaan 120, 3133 AT Vlaardingen, The Netherlands*

(P8) BIOPROCESSING OF INTERFERON ALPHA 2B USING IONIC-LIQUID-BASED SYSTEMS

Augusto Q. Pedro^{1*}, Joana Antunes¹, Patrícia Pereira², Luís A. Passarinha³, Fani Sousa³, João A. P. Coutinho¹, Mara G. Freire¹